

FACT: COVID-19 vaccines will not give you COVID-19

None of the COVID-19 vaccines currently in development in the United States use the live virus that causes COVID-19. There are several different types of vaccines in development. However, the goal for each of them is to teach our immune systems how to recognize and fight the virus that causes COVID-19. Sometimes this process can cause symptoms, such as fever. These symptoms are normal and are a sign that the body is building immunity. Learn more about how COVID-19 vaccines work.

It typically takes a few weeks for the body to build immunity after vaccination. That means it's possible a person could be infected with the virus that causes COVID-19 just before or just after vaccination and get sick. This is because the vaccine has not had enough time to provide protection.

FACT: COVID-19 vaccines will not cause you to test positive on COVID-19 viral tests

Vaccines currently in clinical trials in the United States won't cause you to test positive on viral tests, which are used to see if you have a current infection.

If your body develops an immune response, which is the goal of vaccination, there is a possibility you may test positive on some antibody tests. Antibody tests indicate you had a previous infection and that you may have some level of protection against the virus. Experts are currently looking at how COVID-19 vaccination may affect antibody testing results.

FACT: Getting vaccinated can help prevent getting sick with COVID-19

While many people with COVID-19 have only a mild illness, others may get a [severe illness](#) or they may even die. There is no way to know how COVID-19 will affect you, even if you are not at [increased risk of severe complications](#). If you get sick, you also may spread the disease to friends, family, and others around you while you are sick. COVID-19 vaccination helps

protect you by creating an antibody response without having to experience sickness. Learn more about [how COVID-19 vaccines work](#).

FACT: People who have gotten sick with COVID-19 may still benefit from getting vaccinated

Due to the severe health risks associated with COVID-19 and the fact that re-infection with COVID-19 is possible, people may be advised to get a COVID-19 vaccine even if they have been sick with COVID-19 before.

At this time, experts do not know how long someone is protected from getting sick again after recovering from COVID-19. The immunity someone gains from having an infection, called natural immunity, varies from person to person. Some early evidence suggests natural immunity may not last very long.

We won't know how long immunity produced by vaccination lasts until we have a vaccine and more data on how well it works.

Both natural immunity and vaccine-induced immunity are important aspects of COVID-19 that experts are trying to learn more about, and CDC will keep the public informed as new evidence becomes available.

FACT: Receiving an mRNA vaccine will not alter your DNA


mRNA stands for messenger ribonucleic acid and can most easily be described as instructions for how to make a protein or even just a piece of a protein. mRNA is not able to alter or modify a person's genetic makeup (DNA). The mRNA from a COVID-19 vaccine never enter the nucleus of the cell, which is where our DNA are kept. This means the mRNA does not affect or interact with our DNA in any way. Instead, COVID-19 vaccines that use mRNA work with the body's natural defenses to safely develop protection (immunity) to disease. Learn more about [how COVID-19 mRNA vaccines work](#).

Information obtained from the [CDC website](#).



COVID Vaccine Do's and Don'ts

- **DO**: consider an ice pack on your arm if it is sore after the vaccination
- **DON'T**: panic if you do experience side effects
- **DO**: report any significant signs or symptoms after being vaccinated to your healthcare provider or on the V-safe App
- **DO**: keep your vaccination card in a safe place; consider taking a picture of it and store on your phone in case it gets lost
- **DON'T**: schedule a mammogram or other diagnostic imaging test within two weeks after your COVID-19 vaccine dose
- **DO**: plan to wait 14 days between receiving the COVID-19 vaccine and a different vaccine (e.g., the Shingles vaccine)
- **DON'T**: take medication to pre-treat for any possible side-effects (e.g., don't pre-emptively take Tylenol)
- **DO**: take pain relievers, such as ibuprofen, 7 hours after the vaccine if you are uncomfortable



Health**Works**
Building **Healthy** Worksites

- www.cincyhealthworks.com
- 513.751.1288
- info@cincyhealthworks.com
- 4329-B Red Bank Road
Cincinnati, OH 45227